

Toxicology and Carcinogenesis Studies in Haploinsufficient p16^{lnk4a} /p19^{Arf} N2 Mice (B6.129-Cdkn2a^{tm1Rdp})

Glycidol - GMM 13



Haploinsufficient p16^{lnk4a} /p19^{Arf} Mice Glycidol Studies

Glycidol

CAS No. 556-52-5 $C_3H_6O_2$ MW 74.08

- Chemical intermediate
- Stabilizer in the manufacture of vinyl polymers
- Additive for oil and synthetic hydraulic fluids
- IARC Group 2A probable human carcinogen
- NTP ROC reasonably anticipated to be a human carcinogen



Zymbal gland

Thyroid

NTP Studies: Glycidol Multisite Carcinogen

Glycidol TR 374 (oral gavage) - F344 Rat and B6C3F1 Mouse

Male Rat Female Rat Male Mouse Harderian gland Mesothelioma Mammary gland Forestomach Mammary gland Brain Brain Skin Oral mucosa Forestomach Clitoral gland Liver Thyroid Skin Intestine Leukemia Skin

Female Mouse
Harderian gland
Mammary gland
Uterus
Subcutaneous tissue


Negative in male and female p53(+/-) mice

Lung


Glycidol Oral Gavage Study Designs:


- Doses selected to overlap NIEHS/NTP Glycidol study doses
 - NTP 2-year B6C3F1 mouse study
 - 0, 25, 50 mg/kg
 - NIEHS 26-week p53(-/+) mouse study
 - 0, 25, 50 mg/kg
- NTP 40-week haploinsufficient p16^{INK4a}/p19^{Arf} mouse study
 - 0, 25, 50, 100, 200 mg/kg
 - 15 animals/dose/sex

Survival Curves for 40-week Glycidol Study in Haploinsufficient p16^{lnk4a}/p19^{Arf} Mice



Body Weight - Male and Female Mice





Glycidol: Males & Females - Neoplastic Lesions - Histiocytic Sarcomas

Dose (mg/kg)	0	25	50	100	200
Males	2/15** ^a (13%)	6/15 (40%)	9/15** (60%)	5/15 (33%)	11/15** (73%)
Females	9/15 (60%)	9/15 (60%)	12/15 (80%)	10/15 (67%)	13/15 (87%)

^{**} $p \le 0.01$

^aTrend statistic

Glycidol: Males - Neoplastic Lesions - Lung

Dose (mg/kg)	0	25	50	100	200
Alveolar/Bronchiolar	1/15	0/15	2/15	7/15*	3/15
Adenoma	(7%)	(0%)	(13%)	(47%)	(20%)
Alveolar/Bronchiolar	2/15	1/15	1/15	3/15	0/15
Carcinoma	(13%)	(7%)	(7%)	(20%)	(0%)
Alveolar/Bronchiolar	3/15	1/15	3/15	8/15	3/15
Adenoma or Carcinoma	(20%)	(7%)	(20%)	(53%)	(20%)

^{*} $p \le 0.05$

Glycidol: Females - Neoplastic Lesions - Lung

Dose (mg/kg)	0	25	50	100	200
Alveolar/Bronchiolar	0/15** ^a	1/15	0/15	1/15	4/15*
Adenoma	(0%)	(7%)	(0%)	(7%)	(27%)
Alveolar/Bronchiolar	0/15	0/15	0/15	1/15	0/15
Carcinoma	(0%)	(0%)	(0%)	(7%)	(0%)
Alveolar/Bronchiolar	0/15** ^a	1/15	0/15	2/15	4/15*
Adenoma or Carcinoma	(0%)	(7%)	(0%)	(13%)	(27%)

 $p \le 0.05$ $p \le 0.01$

^aTrend statistic

Glycidol: Males and Females - Nonneoplastic and Neoplastic Lesions of the Forestomach

Dose (mg/kg)	0	25	50	100	200
Males: epithelium hyperplasia	0/15** ^a	1/15	1/15	0/15	6/15**
	(0%)	(7%)	(7%)	(0%)	(40%)
Males: squamous papilloma	0/15	0/15	0/15	0/15	1/15
	(0%)	(0%)	(0%)	(0%)	(7%)
Females: epithelium	0/15	0/15	0/15	1/15	4/15*
Hyperplasia	(0%)	(0%)	(0%)	(7%)	(27%)
Females: squamous papilloma	0/15**a	0/15	0/15	1/15	3/15
	(0%)	(0%)	(0%)	(7%)	(20%)

^{*} $p \le 0.05$ ** $p \le 0.01$

^aTrend statistic

Glycidol: Males - Non-neoplastic Brain Lesions

Dose (mg/kg)	0	25	50	100	200
Neuronopathy	0/15	0/15	0/15	0/15	5/15*
Gliosis	0/15	0/15	0/15	0/15	4/15*
Hemorrhage	0/15	0/15	0/15	0/15	2/15


^{*} $p \le 0.05$

Glycidol: Females - Non-neoplastic Brain Lesions

Dose (mg/kg)	0	25	50	100	200
Neuronopathy	0/15	0/15	0/15	1/15	4/15*
Gliosis	0/15	0/15	0/15	0/15	4/15*
Hemorrhage	0/15	0/15	0/15	1/15	0/15


^{*} $p \le 0.05$

Neuronopathy - symmetrical vacuolization, midbrain nuclei



Thalamus

Neuronopathy - vacuolization of neurons and glial cells



Glycidol Conclusions

- Clear evidence of carcinogenic activity in male haploinsufficient p16^{Ink4a} /p19^{Arf} mice
 - Histiocytic sarcomas
 - Alveolar/bronchiolar adenomas
- Some evidence of carcinogenic activity in female haploinsufficient p16^{lnk4a} /p19^{Arf} mice
 - Alveolar/bronchiolar adenoma
 - Forestomach papillomas
- Treatment-related forestomach and brain lesions in male and female mice
 - Forestomach hyperplasia
 - Neuronopathy